

1 **probit: Probit Regression for Dichotomous Dependent Variables**

Use probit regression to model binary dependent variables specified as a function of a set of explanatory variables.

1.0.1 Syntax

```
> z.out <- zelig(Y ~ X1 + X2, model = "probit", data = mydata)
> x.out <- setx(z.out)
> s.out <- sim(z.out, x = x.out, x1 = NULL)
```

1.0.2 Additional Inputs

In addition to the standard inputs, `zelig()` takes the following additional options for probit regression:

- **robust**: defaults to `FALSE`. If `TRUE` is selected, `zelig()` computes robust standard errors via the `sandwich` package (see [7]). The default type of robust standard error is heteroskedastic and autocorrelation consistent (HAC), and assumes that observations are ordered by time index.

In addition, **robust** may be a list with the following options:

- **method**: Choose from
 - * `"vcovHAC"`: (default if **robust** = `TRUE`) HAC standard errors.
 - * `"kernHAC"`: HAC standard errors using the weights given in [1].
 - * `"weave"`: HAC standard errors using the weights given in [4].
- **order.by**: defaults to `NULL` (the observations are chronologically ordered as in the original data). Optionally, you may specify a vector of weights (either as **order.by** = `z`, where `z` exists outside the data frame; or as **order.by** = `~z`, where `z` is a variable in the data frame). The observations are chronologically ordered by the size of `z`.
- **...**: additional options passed to the functions specified in **method**. See the `sandwich` library and [7] for more options.

1.0.3 Examples

Attach the sample turnout dataset:

```
> data(turnout)
```

Estimate parameter values for the probit regression:

```
> z.out <- zelig(vote ~ race + educate, model = "probit", data = turnout)
> summary(z.out)
```

Set values for the explanatory variables to their default values.

```
> x.out <- setx(z.out)
```

Simulate quantities of interest from the posterior distribution.

```
> s.out <- sim(z.out, x = x.out)
```

```
> summary(s.out)
```

1.0.4 Model

Let Y_i be the observed binary dependent variable for observation i which takes the value of either 0 or 1.

- The *stochastic component* is given by

$$Y_i \sim \text{Bernoulli}(\pi_i),$$

where $\pi_i = \Pr(Y_i = 1)$.

- The *systematic component* is

$$\pi_i = \Phi(x_i\beta)$$

where $\Phi(\mu)$ is the cumulative distribution function of the Normal distribution with mean 0 and unit variance.

1.0.5 Quantities of Interest

- The expected value (`qi$ev`) is a simulation of predicted probability of success

$$E(Y) = \pi_i = \Phi(x_i\beta),$$

given a draw of β from its sampling distribution.

- The predicted value (`qi$pr`) is a draw from a Bernoulli distribution with mean π_i .
- The first difference (`qi$fd`) in expected values is defined as

$$\text{FD} = \Pr(Y = 1 \mid x_1) - \Pr(Y = 1 \mid x).$$

- The risk ratio (`qi$rr`) is defined as

$$\text{RR} = \Pr(Y = 1 \mid x_1) / \Pr(Y = 1 \mid x).$$

- In conditional prediction models, the average expected treatment effect (`att.ev`) for the treatment group is

$$\frac{1}{\sum_{i=1}^n t_i} \sum_{i:t_i=1}^n \{Y_i(t_i = 1) - E[Y_i(t_i = 0)]\},$$

where t_i is a binary explanatory variable defining the treatment ($t_i = 1$) and control ($t_i = 0$) groups. Variation in the simulations are due to uncertainty in simulating $E[Y_i(t_i = 0)]$, the counterfactual expected value of Y_i for observations in the treatment group, under the assumption that everything stays the same except that the treatment indicator is switched to $t_i = 0$.

- In conditional prediction models, the average predicted treatment effect (`att.pr`) for the treatment group is

$$\frac{1}{\sum_{i=1}^n t_i} \sum_{i:t_i=1}^n \{Y_i(t_i = 1) - Y_i(\widehat{t_i = 0})\},$$

where t_i is a binary explanatory variable defining the treatment ($t_i = 1$) and control ($t_i = 0$) groups. Variation in the simulations are due to uncertainty in simulating $Y_i(\widehat{t_i = 0})$, the counterfactual predicted value of Y_i for observations in the treatment group, under the assumption that everything stays the same except that the treatment indicator is switched to $t_i = 0$.

1.0.6 Output Values

The output of each Zelig command contains useful information which you may view. For example, if you run `z.out <- zelig(y ~ x, model = "probit", data)`, then you may examine the available information in `z.out` by using `names(z.out)`, see the `coefficients` by using `z.out$coefficients`, and a default summary of information through `summary(z.out)`. Other elements available through the `$` operator are listed below.

- From the `zelig()` output object `z.out`, you may extract:
 - `coefficients`: parameter estimates for the explanatory variables.
 - `residuals`: the working residuals in the final iteration of the IWLS fit.
 - `fitted.values`: a vector of the in-sample fitted values.
 - `linear.predictors`: a vector of $x_i\beta$.
 - `aic`: Akaike's Information Criterion (minus twice the maximized log-likelihood plus twice the number of coefficients).
 - `df.residual`: the residual degrees of freedom.

- `df.null`: the residual degrees of freedom for the null model.
- `data`: the name of the input data frame.
- From `summary(z.out)`, you may extract:
 - `coefficients`: the parameter estimates with their associated standard errors, p -values, and t -statistics.
 - `cov.scaled`: a $k \times k$ matrix of scaled covariances.
 - `cov.unscaled`: a $k \times k$ matrix of unscaled covariances.
- From the `sim()` output object `s.out`, you may extract quantities of interest arranged as matrices indexed by simulation \times \mathbf{x} -observation (for more than one \mathbf{x} -observation). Available quantities are:
 - `qi$ev`: the simulated expected values, or predicted probabilities, for the specified values of \mathbf{x} .
 - `qi$pr`: the simulated predicted values drawn from the distributions defined by the predicted probabilities.
 - `qi$fd`: the simulated first differences in the predicted probabilities for the values specified in \mathbf{x} and $\mathbf{x1}$.
 - `qi$rr`: the simulated risk ratio for the predicted probabilities simulated from \mathbf{x} and $\mathbf{x1}$.
 - `qi$att.ev`: the simulated average expected treatment effect for the treated from conditional prediction models.
 - `qi$att.pr`: the simulated average predicted treatment effect for the treated from conditional prediction models.

How to Cite the Logit Model

Kosuke Imai, Olivia Lau, and Gary King. *probit: Probit Regression for Dichotomous Dependent Variables*, 2011

How to Cite the Zelig Software Package

To cite Zelig as a whole, please reference these two sources:

Kosuke Imai, Gary King, and Olivia Lau. 2007. “Zelig: Everyone’s Statistical Software,” <http://GKing.harvard.edu/zelig>.

Imai, Kosuke, Gary King, and Olivia Lau. (2008). “Toward A Common Framework for Statistical Analysis and Development.” *Journal of Computational and Graphical Statistics*, Vol. 17, No. 4 (December), pp. 892-913.

See also

The probit model is part of the stats package by **(author?)** [6]. Advanced users may wish to refer to `help(glm)` and `help(family)`, as well as [5]. Robust standard errors are implemented via the sandwich package by **(author?)** [7]. Sample data are from [3].

References

- [1] Donald W.K. Andrews. Heteroskedasticity and autocorrelation consistent covariance matrix estimation. *Econometrica*, 59(3):817–858, May 1991.
- [2] Kosuke Imai, Olivia Lau, and Gary King. *probit: Probit Regression for Dichotomous Dependent Variables*, 2011.
- [3] Gary King, Michael Tomz, and Jason Wittenberg. Making the most of statistical analyses: Improving interpretation and presentation. *American Journal of Political Science*, 44(2):341–355, April 2000. <http://gking.harvard.edu/files/abs/making-abs.shtml>.
- [4] Thomas Lumley and Patrick Heagerty. Weighted empirical adaptive variance estimators for correlated data regression. *jrssb*, 61(2):459–477, 1999.
- [5] Peter McCullagh and James A. Nelder. *Generalized Linear Models*. Number 37 in Monograph on Statistics and Applied Probability. Chapman & Hall, 2nd edition, 1989.
- [6] William N. Venables and Brian D. Ripley. *Modern Applied Statistics with S*. Springer-Verlag, 4th edition, 2002.
- [7] Achim Zeileis. Econometric computing with hc and hac covariance matrix estimators. *Journal of Statistical Software*, 11(10):1–17, 2004.